

References

Research Brief: Anxiety and depression in older adults

- 1) Skoog I. Psychiatric disorders in the elderly. *Can J Psychiatry* 2011 Jul;56(7):387-97.
- 2) Karlsson I, Skoog I. Pyskisk sjukdom hos äldre. *Lakartidningen*. 2013 Nov 6-12;110(45):2000-2.
- 3) American Psychiatric Association. Diagnostic and statistical manual of mental disorders, Fifth edition. Arlington, VA: American Psychiatric Association. 2013.
- 4) American Psychiatric Association. Diagnostic and statistical manual of mental disorders: DSM-IV TR. Washington, DC: American Psychiatric Association. 2000.
- 5) Palsson S, Skoog I. The epidemiology of affective disorders in the elderly: a review. *Int Clin Psychopharmacology* 1997;12 (suppl 2):S3-S13.
- 6) Skoog I. Mental Disorders Associated with Aging. In: Heggenhougen K, Quah S (eds): *International Encyclopedia of Public Health*, Vol 6. San Diego: Academic Press 2008: pp 174-182.
- 7) Veith RC, Raskind MA. Review. The neurobiology of aging: Does it predispose to depression? *Neurobiol Aging* 1988;9:101-117.
- 8) Blazer DG, Hybels CF. Origins of depression in later life. *Psychol Med* 2005;35:11-12.
- 9) Beekman ATF, Deeg DJH, van Tilburg T, et al. Major and minor depression in later life: a study of prevalence and risk factors. *J Aff Dis* 1995;36:65-75.
- 10) Blazer D, Burchett B, Service C, et al. The association of age and depression among the elderly: An epidemiological exploration. *J Gerontol* 1991;46:M210-215.
- 11) Gum AM, King-Kallimanis B, Kohn R. Prevalence of mood, anxiety, and substance-abuse disorders for older Americans in the national comorbidity survey-replication. *Am J Geriatr Psychiatry*. 2009;17:769-81.
- 12) Byers AL, Yaffe K, Covinsky KE, et al. High occurrence of mood and anxiety disorders among older adults: The National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2010 ;67:489-96.
- 13) McDougall FA, Kvaal K, Matthews FE, et al. Prevalence of depression in older people in England and Wales: the MRC CFA Study. *Psychol Med* 2007;37:1787-95.
- 14) Steffens DC, Skoog I, Norton MC, et al. Prevalence of depression and its treatment in an elderly population: the Cache County study. *Arch Gen Psychiatry* 2000;57:601-7.
- 15) Bland RC, Newman SC, Orn H. Prevalence of psychiatric disorders in the elderly in Edmonton. *Acta Psychiatr Scand* 1988; 77(suppl 338):57-63.
- 16) Regier DA, Farmer ME, Rae DS, et al. One-month prevalence of mental disorders in the United States and sociodemographic characteristics: the Epidemiologic Catchment Area Study. *Acta Psychiatr Scand* 1993;88:35-47.
- 17) Blazer D, Swartz M, Woodbury M, et al. Depressive symptoms and depressive diagnoses in a community population. *Arch Gen Psychiatry* 1988;45:1078-1084.
- 18) Lehtinen V, Joukamaa M, Lahtela K, et al. Prevalence of mental disorders among adults in Finland: basic results from the Mini Finland Health Survey. *Acta Psychiatr Scand* 1990;81:418-425.
- 19) Jorm AF. Does old age reduce the risk of anxiety and depression? A review of epidemiological studies across the adult life span. *Psychol Med* 2000;30(1):11-22.
- 20) Börjesson-Hansson A, Waern M, Östling S, Gustafson D, Skoog I. One-month prevalence of mental disorders in a population sample of 95-year-olds. *Am J Geriatr Psychiatry* 2011;19(3):284-91.
- 21) Lindsay J, Briggs K, Murphy E. The Guy's/Age Concern survey. Prevalence rates of cognitive impairment, depression and anxiety in an urban elderly community. *Br J Psychiatry* 1989;155:317-329.
- 22) Kay DWK, Henderson AS, Scott R, et al. Dementia and depression among the elderly living in the Hobart community: the effect of the diagnostic criteria on the prevalence rates. *Psychol Med* 1985;15:771-788.
- 23) Lobo A, Saz P, Marcos G, et al. The prevalence of dementia and depression in the elderly community in a Southern European Population. *Arch Gen Psychiatry* 1995;52:497-506.
- 24) Bowling A. The prevalence of psychiatric morbidity among people age 85 and over living at home. Associations with reported somatic symptoms and with consulting behaviour. *Soc Psychiatry Psychiatr Epidemiol* 1990;25:132-140.
- 25) Gurling DM, Huppert FA, Brayne C, et al. Depressive symptoms in the very elderly. Their prevalence and significance. *Int J Geriatr Psychiatry* 1995;10:497-504.
- 26) Bryant C. Anxiety and depression in old age: challenges in recognition and diagnosis. *Int Psychogeriatrics* 2010; 22(4):11-13.
- 27) Palsson SP, Ostling S, Skoog I. The incidence of first-onset depression in a population followed from the age of 70 to 85. *Psychol Med* 2001;31:1159-1168.
- 28) Norton MC, Skoog I, Toone L, et al. Three year incidence of first-onset depressive syndrome in a population sample of older adults: The Cache County Study. *Am J Geriatr Psychiatry* 2006;14:237-245.

- 29) Krasucki C, Howard R, Mann A. The relationship between anxiety disorders and age. *Int J Geriatr Psychiatry* 1998;13:79-99.
- 30) Manela M, Katona C, Livingston G. How common are the anxiety disorders in old age? *Int J Geriatr Psychiatry* 1996;11:65-70.
- 31) Flint AJ. Epidemiology and comorbidity of anxiety disorders in the elderly. *Am J Psychiatry* 1994;151:640-649.
- 32) Ritchie K, Artero S, Beluche I, et al. Prevalence of DSM-IV psychiatric disorder in the French elderly population. *Br J Psychiatry* 2004;184:147-52.
- 33) Weisberg RB. Overview of generalized anxiety disorder: epidemiology, presentation, and course. *J Clin Psychiatry* 2009;70 (Suppl 2):4-9
- 34) Angst J, Gamma A, Baldwin DS, et al. The generalized anxiety spectrum: prevalence, onset, course and outcome. *Eur Arch Psychiatry Clin Neurosci* 2009;259(1):37-45.
- 35) Carter RM, Wittchen HU, Pfister H, et al. One-year prevalence of subthreshold and threshold DSM-IV generalized anxiety disorder in a nationally representative sample. *Depress Anxiety* 2001;13(2):78-88.
- 36) Wittchen HU, Hoyer J. Generalized anxiety disorder: nature and course. *J Clin Psychiatry*. 2001;62 (Suppl 11):15-19; discussion 20-11.
- 37) Hunt C, Issakidis C, Andrews G. DSM-IV generalized anxiety disorder in the Australian National Survey of Mental Health and Well-Being. *Psychol Med* 2002;32(4):649-659.
- 38) Torres AR, Prince MJ, Bebbington PE, et al. Obsessive-compulsive disorder: prevalence, comorbidity, impact, and help-seeking in the British National Psychiatric Morbidity Survey of 2000. *Am J Psychiatry* 2006;163(11):1978-85
- 39) Nestadt G, Bienvenu OJ, Cai G, et al. Incidence of obsessive-compulsive disorder in adults. *J Nerv Ment Dis* 1998; 186(7):401-6
- 40) Beekman AT, Bremner MA, Deeg DJ, et al. Anxiety disorders in later life: a report from the Longitudinal Aging Study Amsterdam. *Int J Geriatr Psychiatry* 1998;13(10):717-726.
- 41) Chou KL. Specific phobia in older adults: evidence from the national epidemiologic survey on alcohol and related conditions. *Am J Geriatr Psychiatry* 2009;17(5): 376-386.
- 42) Kessler RC, Chiu WT, Demler O, et al. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005;62(6):617-627.
- 43) Stinson FS, Dawson DA, Patricia Chou S, et al. The epidemiology of DSM-IV specific phobia in the USA: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychol Med* 2007;37(7):1047-1059.
- 44) Sigström R. Psychiatric symptoms and disorders in old age: prevalence, course, and diagnostic thresholds (dissertation). University of Gothenburg 2015.
- 45) Liddell A, Locker D, Burman D. Self-reported fears (FSS-II) of subjects aged 50 years and over. *Behaviour Research and Therapy* 1991;29(2):105-112.
- 46) Kirmizioglu Y, Dogan O, Kugu N, et al. Prevalence of anxiety disorders among elderly people. *Int J Geriatr Psychiatry* 2009;24(9):1026-1033.
- 47) Boyd JH, Rae DS, Thompson JW, et al. Phobia: prevalence and risk factors. *Soc Psychiatry Psychiatr Epidemiol* 1990; 5(6):314-323.
- 48) Lindesay J. Phobic disorders in the elderly. *Br J Psychiatry* 1991;159:531-541.
- 49) Sigström R, Ostling S, Karlsson B, Waern M, Gustafson D, Skoog I. A population-based study on phobic fears and DSM-IV specific phobia in 70-year olds. *J Anxiety Disord*. 2011;25(1):148-53.
- 50) Cairney J, McCabe L, Veldhuizen S, et al. Epidemiology of social phobia in later life. *Am J Geriatr Psychiatry*. 2007;15(3):224-33.
- 51) Mohammadi MR, Ghanizadeh A, Mohammadi M, et al. Prevalence of social phobia and its comorbidity with psychiatric disorders in Iran. *Depress Anxiety* 2006;23(7):405-11.
- 52) Wells JE, Browne MA, Scott KM, et al. Prevalence, interference with life and severity of 12 month DSM-IV disorders in Te Rau Hinengaro: the New Zealand Mental Health Survey. *Aust N Z J Psychiatry* 2006;40(10):845-54.
- 53) Trollor JN, Anderson TM, Sachdev PS, et al. Prevalence of mental disorders in the elderly: the Australian National Mental Health and Well-Being Survey. *Am J Geriatr Psychiatry* 2007;15(6):455-66.
- 54) Prévaille M, Boyer R, Grenier S, et al. The epidemiology of psychiatric disorders in Quebec's older adult population. *Can J Psychiatry*. 2008;53:822-32.
- 55) Ernst C and Angst J. Depression in old age. Is there a real decrease in prevalence? A review. *European Archives of Psychiatry and Clinical Neuroscience* 1995;245: 272 -287.
- 56) Skoog I, Nilsson L, Landahl S, Steen B. Mental disorders and the use of psychotropic drugs in an 85-year-old urban population. *Int Psychogeriatrics* 1993;5:33-48
- 57) Baiyewu O, Smith Gamble V, Lane KA, et al. Prevalence estimates of depression in elderly community-dwelling African Americans in Indianapolis and Yoruba in Ibadan, Nigeria. *Int Psychogeriatrics* 2007;19:679-689.
- 58) Beekman ATF, Copeland JR, Prince MJ. Review of community prevalence of depression in later life. *Br J Psychiatry* 1999;174:307 - 311.
- 59) Castro-Costa E, Dewey M, Stewart R, et al. Prevalence of depressive symptoms and syndromes in later life in ten European countries: the SHARE study. *Br J Psychiatry*. 2007 Nov;191:393-401
- 60) Larkin BA, Copeland JR, Dewey ME, et al. The natural history of neurotic disorder in an elderly urban population. Findings from the Liverpool longitudinal study of continuing health in the community. *Br J Psychiatry* 1992;160:681-686.

- 61) Nilsson J, Ostling S, Waern M, Karlsson B, Sigström R, Guo X, Skoog I. The 1-month prevalence of Generalized Anxiety Disorder according to DSM-IV, DSM-V and ICD-10 among non-demented 75-year-olds in Gothenburg, Sweden. *Am J Geriatr Psychiatry*. 2012;20:963-72.
- 62) Bebbington PE. Epidemiology of obsessive-compulsive disorder. *Br J Psychiatry Suppl* 1998(35):2-6
- 63) Grenier S, Préville M, Boyer R, et al. Prevalence and correlates of obsessive-compulsive disorder among older adults living in the community. *J Anxiety Disord* 2009;23:858-65.
- 64) Fredén Klenfeldt I, Karlsson B, Sigström R, Waern M, Östling S, Marlow T, Gustafson D, Skoog I. Prevalence of Obsessive-compulsive Disorder in Relation to Depression and Cognition in an Elderly Population. *Am J Geriatr Psychiatry*. 2014 Mar;22(3):301-8.
- 65) Schneier FR, Johnson J, Hornig CD, et al. Social phobia. Comorbidity and morbidity in an epidemiologic sample. *Arch Gen Psychiatry*. 1992;49(4):282-8.
- 66) Karlsson B, Klenfeldt IF, Sigström R, et al. Prevalence of social phobia in non-demented elderly from a Swedish population study. *Am J Geriatr Psychiatry* 2009;17:127-35.
- 67) Luijendijk HJ, van den Berg JF, Dekker MJHJ, et al. Incidence and recurrence of late-life depression. *Arch Gen Psychiatry* 2008;65:1394-1401.
- 68) Eaton WW, Kramer M, Anthony JC, et al The incidence of specific DIS/DSM-III mental disorders: Data from the NIMH Epidemiologic Catchment Area Program. *Acta Psychiatr Scand* 1989;79:163-178.
- 69) Copeland JRM, Davidson IA, Dewey ME, et al. Alzheimer's disease, other dementias, depression and pseudodementia: Prevalence, incidence and three-year outcome in Liverpool. *Br J Psychiatry* 1992;161:230-239.
- 70) Blanchard MR, Waterreus A, Mann AH. The nature of depression among older people in Inner London, and the contact with primary care. *Br J Psychiatry* 1994;164:396-402.
- 71) Cairney J, Corna LM, Veldhuizen S, et al. Comorbid depression and anxiety in later life: patterns of association, subjective well-being, and impairment. *Am J Geriatr Psychiatry* 2008;16(3):201-8.
- 72) McCabe L, Cairney J, Veldhuizen S, et al. Prevalence and correlates of agoraphobia in older adults. *Am J Geriatr Psychiatry* 2006;14(6):515-22.
- 73) King-Kallimanis B, Gum AM, Kohn R. Comorbidity of depressive and anxiety disorders for older Americans in the national comorbidity survey-replication. *Am J Geriatr Psychiatry*. 2009;17:782-92
- 74) Kvaal K, McDougall FA, Brayne C, et al. Co-occurrence of anxiety and depressive disorders in a community sample of older people: results from the MRC CFAS (Medical Research Council Cognitive Function and Ageing Study). *Int J Geriatr Psychiatry*. 2008;23:229-37.
- 75) Mohammadi MR, Ghanizadeh A, Rahgozar M, et al. Prevalence of obsessive-compulsive disorder in Iran. *BMC Psychiatry* 2004; 4:2
- 76) Wolitzky-Taylor KB, Castriotta N, Lenze EJ, et al. Anxiety disorders in older adults: a comprehensive review. *Depress Anxiety* 2010;27(2):190-211.
- 77) Kay DWK, Beamish P, Roth M. Old age mental disorders in Newcastle upon Tyne. Part I: A study of prevalence. *Br J Psychiatry* 1964;110:146-158.
- 78) Brown TA, Chorpita BF, Barlow DH. Structural relationships among dimensions of the DSM-IV anxiety and mood disorders and dimensions of negative affect, positive affect, and autonomic arousal. *J Abnorm Psychol* 1998;107(2):179-192.
- 79) Fuhrer R, Antonucci TC, Gagnon M, et al. Depressive symptomatology and cognitive functioning: an epidemiological survey in an elderly community in France. *Psychol Med* 1992; 22:159-172.
- 80) Palsson S, Aevansson O, Skoog I. Depression, cerebral atrophy, cognitive performance and incidence of dementia. Population study of 85-year-olds. *Brit J Psychiatr* 1999;174:249-253.
- 81) Palsson S, Larsson L, Tengelin E, et al. The prevalence of depression in relation to cerebral atrophy and cognitive performance in 70- and 74-year-old women in Gothenburg. The Women's Health Study. *Psychol Med* 2001;31:39-49.
- 82) Palsson S, Johansson B, Berg S et al. A population study on the influence of depression on neuropsychological functioning in 85-year-olds. *Acta Psychiatr Scand* 2000;101:185-193.
- 83) Fichter MM, Bruce ML, Schröppel H, et al. Cognitive impairment and depression in the oldest old in a German and in U.S. communities. *Eur Arch Psychiatry Clin Neuroscience* 1995;245:319-325.
- 84) Beats BC, Sahakian BJ, Levy R. Cognitive performance in tests sensitive to frontal lobe dysfunction in the elderly depressed. *Psychol Med* 1996;26:591-603.
- 85) van Ojen R, Hooijer C, Bezemer D, et al. Late-life depressive disorder in the community. I. The relationship between MMSE score and depression in subjects with and without psychiatric history. *Br J Psychiatry* 1995;166:311-315.
- 86) Mantella RC, Butters MA, Dew MA, et al. Cognitive impairment in late-life generalized anxiety disorder. *Am J Geriatr Psychiatry* 2007;15(8):673-9.
- 87) Jorm AF, van Duijn VM, Chandra V, et al. Psychiatric history and related exposures as risk factors for Alzheimer's disease: a collaborative re-analysis of case-control studies. *Int J Epidemiol* 1991;20(suppl 2):S43-S47.
- 88) Devanand DP, Sano M, Tang MX, et al. Depressed mood and the incidence of Alzheimer's disease in the elderly living in the community. *Arch Gen Psychiatry* 1996;53:175-182.
- 89) Ownby RL, Crocco E, Acevedo A et al. Depression and risk for Alzheimer disease: systematic review, meta-analysis, and metaregression analysis. *Arch Gen Psychiatry* 2006;63:530-538.

- 90) Geerlings MI, den Heijer T, Koudstaal PJ, et al. History of depression, depressive symptoms, and medial temporal lobe atrophy and the risk of Alzheimer disease. *Neurology* 2008;70(15):1258-64.
- 91) Dotson VM, Beydoun MA, Zonderman AB. Recurrent depressive symptoms and the incidence of dementia and mild cognitive impairment. *Neurology* 2010;75(1):27-34.
- 92) Saczynski JS, Beiser A, Seshadri S, et al. Depressive symptoms and risk of dementia: the Framingham Heart Study. *Neurology* 2010;75(1):35-41.
- 93) Becker JT, Chang YF, Lopez OL, et al. Depressed mood is not a risk factor for incident dementia in a community-based cohort. *Am J Geriatr Psychiatry* 2009;17(8):653-63.
- 94) Persson G, Skoog I. Subclinical dementia: Relevance of cognitive and affective symptoms and signs. A nine year prospective study. *J Geriatr Psychiatry Neurol* 1992;5:172-178.
- 95) Henderson AS, Korten AE, Jacomb PA, et al. The course of depression in the elderly: a longitudinal community-based study in Australia. *Psychol Med* 1997;27:119-129.
- 96) Palsson S. Population Studies On Depression in The Elderly. Prevalence, incidence and relation to cognitive function and dementia. Dissertation. University of Gothenburg 2000.
- 97) Mirza SS, de Bruijn RF, Direk N, Hofman A, Koudstaal PJ, Ikram MA, Tiemeier H. Depressive symptoms predict incident dementia during short- but not long-term follow-up period. *Alzheimers Dement*. 2014 Oct;10(5 Suppl):S323-S329.
- 98) Gallacher J, Bayer A, Fish M, et al. Does anxiety affect risk of dementia? Findings from the Caerphilly Prospective Study. *Psychosom Med* 2009;71(6):659-66.
- 99) de Bruijn RF, Direk N, Mirza SS, Hofman A, Koudstaal PJ, Tiemeier H, Ikram MA. Anxiety is not associated with the risk of dementia or cognitive decline: the Rotterdam Study. *Am J Geriatr Psychiatry*. 2014 Dec;22(12):1382-90. doi: 10.1016/j.jagp.2014.03.001.
- 100) Djernes JK. Prevalence and predictors of depression in populations of elderly: a review. *Acta Psychiatr Scand* 2006;113:372-387.
- 101) Bruce ML. Psychosocial risk factors for depressive disorders in late life. *Biol Psychiatry* 2002;52:175 - 184.
- 102) Rihmer Z, Pestalicy P, Pihlgren H, Rutz W. 'Anxiety/aggression-driven depression' and 'male depressive syndrome': are they the same? *Psychiatry Res* 1998;77(3):209-10.
- 103) Socialstyrelsen 2016. Statistik om dödsorsaker 2015. <http://www.socialstyrelsen.se/publikationer2016/2016-8-3>
- 104) Clayton PJ, Halikas JA, Maurice WL. The depression of widowhood. *Br J Psychiatry* 1972;120:71-78.
- 105) Harlow SD, Goldberg EL, Comstock GW. A longitudinal study of the prevalence of depressive symptomatology in elderly widowed and married woman. *Arch Gen Psychiatry* 1991;48:1065-1068.
- 106) Geerlings SW, Beekman AT, Deeg DJ et al. Physical health and the onset and persistence of depression in older adults: an eight-wave prospective community-based study. *Psychol Med* 2000;30:369-380.
- 107) Beekman ATF, Kriegsman DMW, Deeg DJH, et al. The association of physical health and depressive symptoms in the older population: age and sex differences. *Social Psychiatry and Psychiatric Epidemiology* 1995;30:32-38.
- 108) Braam AW, Prince MJ, Beekman ATF, et al. Physical health and depressive symptoms in older Europeans: results from EURODEP. *Brit J Psychiatry* 2005;187:35-42.
- 109) Gureje O, Ademola A, Olley BO. Depression and disability: comparisons with common physical conditions in the Ibadan Study of Aging. *J Am Geriatr Soc* 2008;56:2033-2038.
- 110) Prince MJ, Harwood R, Thomas A, et al. Gospel Oak V. Impairment, disability and handicap as risk factors for depression in old age. The Gospel Oak Project V. *Psychol Med* 1997;27:311-321.
- 111) Braam AW, Delespaul P, Beekman ATF, et al. National context of healthcare, economy and religion and the association between disability and depressive symptoms in older Europeans: results from the EURODEP concerted action. *Eur J Ageing* 2004;1:26-36.
- 112) Weyerer S, Häfner H, Mann AH, et al. Prevalence and course of depression among elderly residential home admissions in Mannheim and Camden, London. *Int Psychogeriatrics* 1995;7:479-493.
- 113) Henderson AS, Jorm AF, Mackinnon A, et al. The prevalence of depressive disorders and the distribution of depressive symptoms in later life: a survey using Draft ICD-10 and DSM-III-R. *Psychol Med* 1993;23:719-729.
- 114) Dhondt ADF, Hooijer C. Iatrogenic origins of depression in the elderly. Is medication a significant etiologic factor in geriatric depression? Considerations and preliminary approach. *Int J Geriatric Psychiatry* 1995;10:1-8.
- 115) Green BH, Copeland JRM, Dewey ME, et al. Risk factors for depression in elderly people: a prospective study. *Acta Psychiatr Scand* 1992;86:213-217.
- 116) Prince MJ, Harwood R, Thomas A, et al. Social support deficits, loneliness and life events as risk factors for depression in old age. The Gospel Oak Project VI. *Psychol Med* 1997;27:323-332.
- 117) Areán PA, Reynolds CFI. The impact of psychosocial factors on late-life depression. *Biol Psychiatry* 2005;58:277-282.
- 118) Skoog I. Neuropsychiatric aspects of vascular cognitive impairment. In: Miyoshi K, Morimura Y, Maeda K, editors. *Neuropsychiatric Disorders*. Springer Japan KK, Tokyo 2010: 149-162.
- 119) van Ojen R, Hooijer C, Jonker C, et al. Late-life depressive disorder in the community, early onset and the decrease of vulnerability with increasing age. *J Aff Dis* 1995;33:159-166.
- 120) Cederblad M, Dahlin L, Hagnell O, Hansson K. Intelligence and temperament as protective factors for mental health. A cross-sectional and prospective epidemiological study. *Eur Arch Psychiatry Clin Neurosci* 1995;245:11-19.

- 121) Duberstein PR, Palsson SP, Waern M, et al. Personality and risk for depression in a birth cohort of 70-year-olds followed for 15 years. *Psychol Med* 2008;38:663-71
- 122) Bekaroglu M, Uluuku N, Tanriöver S, et al. Depression in an elderly population in Turkey. *Acta Psychiatr Scand* 1991;84:174-178.
- 123) Gudmundsson P, Andersson S, Gustafson D, Waern M, Ostling S, Hällström T, Palsson S, Skoog I, Hulthen L. Depression in Swedish women: relationship to factors at birth. *Eur J Epidemiol* 2011;26(1):55-60.
- 124) Gudmundsson P, Lindwall M, Gustafson DR, Östling S, Hällström T, Waern M, Ingmar Skoog I. Longitudinal associations between physical activity and depression scores in women followed 32 years. *Acta Psych Scand* 2015 Dec;132(6):451-8..
- 125) Vink D, Aartsen MJ, Comijs HC, Heymans MW, Penninx BW, Stek ML, Deeg DJ, Beekman AT. Onset of anxiety and depression in the aging population: comparison of risk factors in a 9-year prospective study. *Am J Geriatr Psychiatry* 2009;17:642-652.
- 126) Corna LM, Cairney J, Herrmann N, Veldhuizen S, McCabe L, Streiner D. Panic disorder in later life: results from a national survey of Canadians. *Int Psychogeriatr*. 2007;19(6):1084-96.
- 127) Vink D, Aartsen MJ, Schoevers RA. Risk factors for anxiety and depression in the elderly: a review. *J Affect Disord*. 2008;106(1-2):29-44.
- 128) Simoni M, Pantoni L, Pracucci G, et al. Prevalence of CT-detected cerebral abnormalities in an elderly Swedish population sample. *Acta Neurol Scand* 2008;118:260-267.
- 129) Beats B, Levy R, Förstl H. Ventricular enlargement and caudate hyperdensity in elderly depressives. *Biol Psychiatry* 1991;30:452-458.
- 130) Alexopoulos GS, Young RC, Shindledecker RD. Brain computed tomography findings in geriatric depression and primary degenerative dementia. *Biol Psychiatry* 1992;31:591-599.
- 131) Elkins H, Friedman L, Wise A, et al. Meta-analysis of studies of ventricular enlargement and cortical sulcal prominence in mood disorders. Comparisons with controls or patients with schizophrenia. *Arch Gen Psychiatry* 1995;52:735-746.
- 133) Rabins PV, Pearlson GD, Aylward E, et al. Cortical magnetic resonance imaging changes in elderly inpatients with major depression. *Am J Psychiatry* 1991;148:617-620.
- 133) Schweitzer I, Tuckwell V, Ames D, et al. Structural neuroimaging studies in late-life depression: a review. *World J Biol Psychiatry* 2001;2:83-88.
- 134) Krishnan KR, McDonald WM, Doraiswamy PM, et al. Neuroanatomical substrates of depression in the elderly. *Eur Arch Psychiatry Clin Neuroscience* 1993;243:41-46.
- 135) Rainer MK, Mucke HA, Zehetmayer S, et al. Data from the VITA Study do not support the concept of vascular depression. *Am J Geriatr Psychiatry* 2006;14:531-537.
- 136) Almeida OP, Burton EJ, Ferrier N, et al. Depression with late onset is associated with right frontal lobe atrophy. *Psychol Med* 2003;33:675-681.
- 137) Kumar A, Bilker W, Jin Z, et al. Atrophy and high intensity lesions: complementary neurobiological mechanisms in late-life major depression. *Neuropsychopharmacology* 2000;22:264-274.
- 138) Kumar A, Mintz J, Bilker W, et al. Autonomous neurobiological pathways to late-life major depressive disorder: clinical and pathophysiological implications. *Neuropsychopharmacology* 2002;26:229-236.
- 139) Olesen PJ, Gustafson DR, Simoni M, Pantoni L, Ostling S, Guo X, Skoog I. Temporal Lobe Atrophy and White Matter Lesions are Related to Major Depression over 5 years in the Elderly. *Neuropsychopharmacology* 2010;35(13):2638-45.
- 140) Ikram MA, Luijckendijk HJ, Vernooij MW, et al. Vascular brain disease and depression in the elderly. *Epidemiology* 2010;21:78-81.
- 141) Skoog I, Waern M, Duberstein P, Blennow K, Zetterberg H, Börjesson-Hanson A, Östling S, Guo X, Kern J, Gustafson D, Gudmundsson P, Marlow T, Kern S. A 9-year prospective population-based study on the association between the APOE 4 allele and late-life depression in Sweden. *Biol Psychiatry* 2015 Nov 15;78(10):730-6.
- 142) Liebetrau M, Steen B, Skoog I. Stroke in 85-year-olds. Prevalence, incidence, risk factors and relation to mortality and dementia. *Stroke* 2003;34:2617-2622
- 143) Alexopoulos GS, Meyers BS, Young RC, et al (1997). 'Vascular depression' hypothesis. *Arch Gen Psychiatry* 1997;54: 915-922.
- 144) Cumming TB, Churilov L, Skoog I, et al. Little evidence for different phenomenology in post-stroke depression. *Acta Psychiatr Scand* 2010;121:424-430
- 145) Beblo T, Driessen M. No Melancholia in Poststroke Depression? A Phenomenologic Comparison of Primary and Poststroke Depression. *J Geriatr Psychiatry Neurol* 2002; 15:44-49.
- 146) Hackett ML, Yapa C, Parag V et al. Frequency of depression after stroke: a systematic review of observational studies. *Stroke* 2005;36:1330-40.
- 147) Hackett ML, Anderson CS. Predictors of depression after stroke: a systematic review of observational studies. *Stroke* 2005;36:2296-301.
- 148) Rao R, Jackson S, Howard R. Depression in older people with mild stroke, carotid stenosis and peripheral vascular disease: a comparison with healthy controls. *Int J Geriatr Psychiatry* 2001;16:175-83
- 149) Pohjasvaara T, Leppavuori A, Siira I et al. Frequency and clinical determinants of poststroke depression. *Stroke* 1998;29:2311-7.
- 150) Linden L, Blomstrand C, Skoog I. Depressive disorders after 20 months in elderly stroke patients. A case-control study. *Stroke* 2007;38:1860-1863.
- 151) Robinson RG, Jorge RE. Post-Stroke Depression: A Review. *Am J Psychiatry*. 2016 Mar 1;173(3):221-31.
- 152) Pantoni L, Garcia JH. Pathogenesis of leukoaraiosis: a review. *Stroke* 1997;28: 652-659

- 153) Fernando MS, Ince PG, on behalf of the MRC Cognitive function and Ageing Neuropathology Study Group. Vascular pathologies and cognition in a population-based cohort of elderly people. *J Neurol Sciences* 2004;226:13-17.
- 154) O'Brien J, Desmond P, Ames D, et al. A magnetic resonance imaging study of white matter lesions in depression and Alzheimer's disease. *Br J Psychiatry* 1996;168:477-485.
- 155) Herrmann LL, Le Masurier M, Ebmeier KP. White matter hyperintensities in late life depression: a systematic review. *J Neurol Neurosurg Psychiatry* 2008;79:619-24.
- 156) Godin O, Dufouil C, Maillard P, et al. White matter lesions as a predictor of depression in the elderly: the 3C-Dijon study. *Biol Psychiatry* 2008;63:663-669.
- 157) Steffens DC, Krishnan KR, Crump C, et al. Cerebrovascular disease and evolution of depressive symptoms in the cardiovascular health study. *Stroke* 2002;33:1636-1644.
- 158) Teodorczuk A, O'Brien JT, Firbank MJ, et al. White matter changes and late-life depressive symptoms: longitudinal study. *Br J Psychiatry* 2007;191:212-217.
- 159) Versluis CE, van der Mast RC, van Buchem MA, et al. Progression of cerebral white matter lesions is not associated with development of depressive symptoms in elderly subjects at risk of cardiovascular disease: The PROSPER Study. *Int J Geriatr Psychiatry* 2006;21:375-381.
- 160) O'Brien J, Ames D, Chiu E et al. Severe deep white matter lesions and outcome in elderly patients with major depressive disorder: follow up study. *BMJ* 1998;317:982-984
- 161) Simpson S, Baldwin RC, Jackson A et al. Is subcortical disease associated with a poor response to antidepressants? Neurological, neuropsychological and neuroradiological findings in late-life depression. *Psychol Med* 1998;28:1015-26
- 162) Liebetrau M, Steen B, Skoog I. Depression as a risk factor for the incidence of first-ever stroke in 85-year-olds. *Stroke* 2008;39:1960-5.
- 163) Magruder KM, Calderone GE. Public health consequences of different thresholds for the diagnosis of mental disorders. *Compr Psychiatry* 2000;41(2 Suppl 1):14-8.
- 164) Rowe SK, Rapaport MH. Classification and treatment of sub-threshold depression. *Curr Opin Psychiatry* 2006;19(1):9-13.
- 165) Gurland B. The impact of depression on quality of life of the elderly. *Clin Geriatr Med* 1992;8:377-386.
- 166) Gureje O, Kola L, Afolabi E. Epidemiology of major depressive disorder in elderly Nigerians in the Ibadan Study of Ageing: a community-based survey. *Lancet* 2007;370:957-964.
- 167) Beekman ATF. Depression in later life: Studies in the community (Dissertation). CopyPrint 2000, Enschede 1996.
- 168) Kennedy GJ, Kelman HR, Thomas C, et al. Hierarchy of characteristics associated with depressive symptoms in an urban elderly sample. *Am J Psychiatry* 1989;46:220-225.
- 169) Vaillant GE, Orav J, Meyer SE, et al (1996) Late-life consequences of affective spectrum disorder. *Int Psychogeriatrics* 1996;8:13-32.
- 170) Waern M, Runeson BS, Allebeck P, Beskow J, et al. Mental disorder in elderly suicides: a case-control study. *Am J Psychiatry* 2002;159:450-5
- 171) Saz P, Dewey ME. Depression, depressive symptoms and mortality in persons aged 65 and over living in the community: a systematic review of the literature. *Int J Geriatr Psychiatry* 2001;16:622-30.
- 172) O'Brien JT, Ames D. Why do the depressed elderly die? *Int J Geriatr Psychiatry* 1994;9:689-693.
- 173) Murphy JM, Monson RR, Olivier DC, et al. Affective disorders and mortality. *Arch Gen Psychiatry* 1987;44:473-480.
- 174) Mykletun A, Bjerkeset O, Overland S, et al. Levels of anxiety and depression as predictors of mortality: the HUNT study. *Br J Psychiatr* 2009;195(2):118-25.
- 175) Pennix BWJH, Geerlings SW, Deeg DJH, et al. Minor and major depression and the risk of death in older persons. *Arch Gen Psychiatry* 1999;56:899 - 895.
- 176) Bos MJ, Lindén T, Koudstaal PJ et al. Depressive Symptoms and Risk of Stroke: The Rotterdam Study. *J Neurol Neurosurg Psychiatry* 2008;79:997-1001.
- 177) Salaycik KJ, Kelly-Hayes M, Beiser A et al. Depressive symptoms and risk of stroke: the Framingham Study. *Stroke* 2007;38:16-21.
- 178) Ramasubbu R, Robinson RG, Flint AJ et al (1998) Functional impairment associated with acute poststroke depression: the Stroke Data Bank Study. *J.Neuropsychiatry Clin Neurosci* 1998;10:26-33.
- 179) Schulz R, Beach SR, Ives DG et al. Association between depression and mortality in older adults: the Cardiovascular Health Study. *Arch Intern Med* 2000;160:1761-8.
- 180) Pohjasvaara T, Vataja R, Leppavuori A et al. Depression is an independent predictor of poor long-term functional outcome post-stroke. *Eur J Neurol* 2001;8:315-9.
- 181) Schubert DS, Taylor C, Lee S et al. Detection of depression in the stroke patient. *Psychosomatics* 1992;33:290-4.
- 182) Kotila M, Numminen H, Waltimo O et al. Post-stroke depression and functional recovery in a population-based stroke register. The Finnstroke study. *Eur J Neurol* 1999;6:309-12
- 183) Barth, J., M. Schumacher, and C. Herrmann-Lingen, Depression as a risk factor for mortality in patients with coronary heart disease: a meta-analysis. *Psychosom Med*, 2004. 66(6): p. 802-13.
- 184) Meijer, A., et al., Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis of 25 years of research. *Gen Hosp Psychiatry*. 33(3): p. 203-16.

- 185) van Melle, J.P., et al., Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis. *Psychosom Med*, 2004. 66(6): p. 814-22.
- 186) Jayadevappa, R., et al., The burden of depression in prostate cancer. *Psychooncology*. 21(12): p. 1338-45.
- 187) Ho, P.M., et al., Depression predicts mortality following cardiac valve surgery. *Ann Thorac Surg*, 2005. 79(4): p. 1255-9.
- 188) Lin, E.H., et al., Depression and increased mortality in diabetes: unexpected causes of death. *Ann Fam Med*, 2009. 7(5): p. 414-21.
- 189) Wu Q, Liu J, Gallegos-Orozco JF, Hentz JG. Depression, fracture risk, and bone loss: a meta-analysis of cohort studies. *Osteoporos Int*. 2010 Oct;21(10):1627-35
- 190) Jorge, R.E., et al., Mortality and poststroke depression: a placebo-controlled trial of antidepressants. *Am J Psychiatry*, 2003. 160(10): p. 1823-9. 10.1007/s00198-010-1181-x. Epub 2010 Mar 4)
- 191) Gallo, J.J., et al., The effect of a primary care practice-based depression intervention on mortality in older adults: a randomized trial. *Ann Intern Med*, 2007. 146(10): p. 689-98.
- 192) Bogner, H.R., et al., Diabetes, depression, and death: a randomized controlled trial of a depression treatment program for older adults based in primary care (PROSPECT). *Diabetes Care*, 2007. 30(12): p. 3005-10.
- 193) Gallagher D, Savva GM, Kenny R, Lawlor BA. What predicts persistent depression in older adults across Europe? Utility of clinical and neuropsychological predictors from the SHARE study. *J Affect Disord*. 2013 May;147(1-3):192-7.
- 194) Licht-Strunk E, van der Windt DA, van Marwijk HW, de Haan M, Beekman AT. The prognosis of depression in older patients in general practice and the community. A systematic review. *Fam Pract*. 2007 Apr;24(2):168-80. Epub 2007 Jan 20.
- 195) Beekman AT, Geerlings SW, Deeg DJ, Smit JH, Schoevers RS, de Beurs E, Braam AW, Penninx BW, van Tilburg W. The natural history of late-life depression: a 6-year prospective study in the community. *Arch Gen Psychiatry*. 2002 Jul;59(7):605-11
- 196) Licht-Strunk E, Van Marwijk HW, Hoekstra T, Twisk JW, De Haan M, Beekman AT. Outcome of depression in later life in primary care: longitudinal cohort study with three years' follow-up. *BMJ*. 2009 Feb 23;338:a3079. doi: 10.1136/bmj.a3079.
- 197) Sigström R, Skoog I, Karlsson B, Nilsson J, Östling S. Nine-year follow-up of specific phobia in a population sample of older people. *Depress Anxiety*. 2015 Dec 8. doi: 10.1002/da.22459. [Epub ahead of print]
- 198) Karlsson B, Sigström R, Waern M, Östling S, Gustafson D, Skoog I. The Prognosis and Incidence of Social Phobia in an Elderly Population. A 5-Year Follow-up. *Acta Psych Scand* 2010; 122:4-10
- 199) Saeed Mirza S, Ikram MA, Hofman A, Tiemeier H. Anxiety does not predict mortality. A population-based study. *World Psychiatry*. 2015 Feb;14(1):103-4. doi: 10.1002/wps.20192.
- 200) van Hout HP, Beekman AT, de Beurs E, Comijs H, van Marwijk H, de Haan M, van Tilburg W, Deeg DJ. Anxiety and the risk of death in older men and women. *Br J Psychiatry*. 2004 Nov;185:399-404.
- 201) Carrière I, Ryan J, Norton J, Scali J, Stewart R, Ritchie K, Ancelin ML. Anxiety and mortality risk in community-dwelling elderly people. *Br J Psychiatry*. 2013 Sep;203(3):303-9.
- 202) Vink, D., et al., Onset of anxiety and depression in the aging population: comparison of risk factors in a 9-year prospective study. *Am J Geriatr Psychiatry*, 2009. 17(8): p. 642-52.
- 203) Roy-Byrne, P.P., et al., Anxiety disorders and comorbid medical illness. *Gen Hosp Psychiatry*, 2008. 30(3): p. 208-25.
- 204) Stein MB, Kean YM. Disability and quality of life in social phobia: epidemiologic findings. *Am J Psychiatry* 2000;157(10):1606-13.
- 205) Merrill J, Owens J. Age and attempted suicide. *Acta Psychiatr Scand* 1990;82:385-388.
- 206) Lyness JM, Conwell Y, Nelson JC. Suicide attempts in elderly psychiatric inpatients. *J Am Geriatr Soc* 1992;40:320-324.
- 207) Van Orden KA, et al. Characteristics and comorbid symptoms of older adults reporting death ideation. *Am J Ger Psychiatry*, 2013 Aug;21(8):803-10.
- 208) Skoog I, Aevansson O, Beskow J, Larsson L, Palsson S, Waern M, Landahl S, Östling, S. Suicidal feelings in a population sample of non-demented 85-year-olds. *Am J Psychiatry* 1996;153:1015-1020.
- 209) Wiktorsson S, Runesson B, Skoog I, et al. Attempted suicide in the elderly: Characteristics of suicide attempters (70+) and a general population comparison group. *Am J Geriatr Psychiatry* 2010;18:57-67
- 210) Nowers M. Deliberate self-harm in the elderly: A survey of one London borough. *Int J Geriatr Psychiatry* 1993;8:609-614.
- 211) Draper B. Suicidal behaviour in the elderly. *Int J Geriatr Psychiatry* 1994;9:655-661.
- 212) Conwell Y, Rotenberg M, Caine ED. Completed suicide at age 50 and over. *J Am Geriatr Soc* 1990;38:640-644.
- 213) Waern M. Alcohol dependence and abuse in elderly suicides. *Alcohol and alcoholism* 2003;38:249-254.
- 214) Morin J, et al. Alcohol use disorder in elderly suicide attempters: a comparison study. *Am J Ger Psychiatry* 2013;21(2):196-203.
- 215) Jonson M et al. Anxiety symptoms and suicidal feelings in a population sample of 70-years-olds without dementia. *Int Psychogeriatrics*. 2012 Nov; 24(11):1865-71
- 216) Olsson P, Wiktorsson S, Sacuiu S, Marlow T, Östling S, Fässberg MM, Skoog I, Waern M. Cognitive function in older suicide attempters and a population-based comparison group. *J Geriatr Psychiatry Neurol*. 2016 Jan 25.

- 217) Fässberg MM, Cheung G, Canetto SS, Erlangsen A, Lapierre S, Lindner R, Draper B, Gallo JJ, Wong C, Wu J, Duberstein P, Waern M. A systematic review of physical illness, functional disability, and suicidal behavior among older adults. *Ageing Ment Health*. 2016 Feb;20(2):169-94.
- 218) Mellqvist Fässberg M, Ostling S, Braam AW, Bäckman K, Copeland JR, Fichter M, Kivelä SL, Lawlor BA, Lobo A, Magnússon H, Prince MJ, Reischies FM, Turrina C, Wilson K, Skoog I, Waern M. Functional disability and death wishes in older Europeans: Results from the Eurodep concerted action. *Soc Psychiatry Psychiatr Epidemiol*. 2014 Sept;49(9):1475-82.
- 219) Waern M, Rubenowitz E, Runeson B, Skoog I, Wilhelmson K, Allebeck P. Burden of illness and suicide in elderly people: a case-control study. *Brit Med J* 2002;324:1355-57.
- 220) Wiktorsson S, Berg A, Wilhelmson K, Mellqvist Fässberg M, Van Orden K, Duberstein P, Waern M. Assessing the role of physical illness in young old and old old suicide attempters. *Int J Geriatric Psychiatry*. 2015 Nov 11
- 221) Fässberg MM, van Orden KA, Duberstein P, Erlangsen A, Lapierre S, Bodner E, Canetto SS, De Leo D, Szanto K, Waern M. A systematic review of social factors and suicidal behavior in older adulthood. *Int J Environ Res Public Health*. 2012 Mar;9(3):722-745.
- 222) Fässberg MM, Östling S, Börjesson-Hanson A, Skoog I, Waern M. Suicidal feelings in the twilight of life: a cross-sectional population-based study of 97-year-olds. *BMJ Open*. 2013 Feb 1;3(2).
- 223) Szanto K, Lenze E, Waern M, Duberstein P, Bruce M, Epstein-Lubow G, Conwell Y. Research to reduce the suicide rate in older adults: methodology roadblocks and candidate paradigms. *Psychiatric Services*. 2013 Jun;64(6):586-9.
- 224) Lapierre S, Erlangsen A, Waern M, De Leo D, Oyama H, Scocco P, Gallo J, Szanto K, Conwell Y, Draper B, Quinnett P; International Research Group for Suicide Among the Elderly. A systematic review of elderly suicide prevention programs. *Crisis*. 2011;32(2):88-98.
- 225) von Knorring L, Ekselius L. Idiopathic pain and depression. *Qual Life Res*. 1994;3 Suppl 1:S57-68)
- 226) Thompson T, Correll CU, Gallop K, Vancampfort D, Stubbs B. Pain Perception Altered in People With Depression? A Systematic Review and Meta-Analysis of Experimental Pain Research. *J Pain*. 2016 Aug 31. pii: S1526-5900(16)30202-4.
- 227) Kallioinen M, Bernhardsson J, Grohp M, Lisspers J, Sundin O. [Mental illness a problem among primary health care patients. A questionnaire comprising more than 2 000 patients]. *Lakartidningen*. 2010;107(23):1545-7.
- 228) Lotfi L, Flyckt L, Krakau I, Mårtensson B, Nilsson GH. Undetected depression in primary healthcare: occurrence, severity and co-morbidity in a two-stage procedure of opportunistic screening. *Nordic Journal of Psychiatry*. 2010;64(6):421-7.
- 229) Forsman AKI, Nordmyr J, Wahlbeck K. Psychosocial interventions for the promotion of mental health and the prevention of depression among older adults. *nHealth Promot Int*. 2011 Dec;26 Suppl 1:i85-107. doi: 10.1093/heapro/dar074.
- 230) Djukanović I, Sorjonen K, Peterson U. Association between depressive symptoms and age, sex, loneliness and treatment among older people in Sweden. *Ageing Ment Health*. 2015;19(6):560-8.
- 231) SBU. Behandling av depression hos äldre. En systematisk litteraturoversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2015. SBU-rapport nr 233. ISBN 978-91-85413-76-8.
- 232) Nelson JC, Delucchi K, Schneider LS. Efficacy of second generation antidepressants in late-life depression: a meta-analysis of the evidence. *Am J Geriatr Psychiatry*. 2008 Jul;16(7):558-67. doi: 10.1097/JGP.0b013e3181693288.
- 233) Cuijpers P, van Straten A, Smit F, Andersson G. Is psychotherapy for depression equally effective in younger and older adults? A meta-regression analysis. *International journal of psychogeriatrics*. 2009;11:16-24.
- 234) Spaans HP, Sienaert P, Bouckaert F, van den Berg JF, Verwijk E, Kho KH, Stek ML, Kok RM. Speed of remission in elderly patients with depression: electroconvulsive therapy v. medication. *Br J Psychiatry*. 2015 Jan;206(1):67-71. doi: 10.1192/bjp.bp.114.148213. Epub 2014 Oct 16.
- 235) Nordanskog P, Hultén M, Landén M, Lundberg J, von Knorring L, Nordenskjöld A. Electroconvulsive Therapy in Sweden 2013: Data From the National Quality Register for ECT. *J ECT*. 2015 Dec;31(4):263-7.
- 236) Verwijk E, Comijs HC, Kok RM, Spaans HP, Tielkes CE, Scherder EJ, Stek ML. Short- and long-term neurocognitive functioning after electroconvulsive therapy in depressed elderly: a prospective naturalistic study. *Int Psychogeriatr*. 2014;26:315-24.